

How To Do the Rapid Test for Malaria

Modified for training in the use of the Generic Pan-Pf Test for falciparum and non-falciparum malaria



READ THESE INSTRUCTIONS CAREFULLY BEFORE YOU BEGIN.


1. Check the expiry date on the test packet.



2. Put on the gloves. Use new gloves for each patient.




3. Open the packet and remove:




4. Write the patient's name on the test.



5. Open the alcohol swab. Grasp the 4th finger on the patient's left hand. Clean the finger with the alcohol swab.




6. Open the lancet. Prick patient's finger to get a drop of blood. Do not allow the tip of the lancet to touch anything before pricking the patient's finger.



7. Discard the lancet in the Sharps Box immediately after pricking finger.

c. Desiccant sachet


8. Use the capillary tube to collect the drop of blood.



14. How to read the test results:


POSITIVE

A line near letter "C" followed by ONE OR TWO LINES near letter "T" means the patient is positive for malaria as shown below. (Test is positive even if the test lines are faint.)




NEGATIVE

A line near letter "C" followed by **NO LINES** near letter "**T**" means the patient **DOES NOT** have either falciparum malaria or non-falciparum malaria.




INVALID RESULT

NO LINE near letter "C" and one or two lines or no line near letter "T" means the test is INVALID.



Repeat the test using a new RDT if no control line appears.

If no line appears near the letter "C," repeat the test using a **NEW unopened** test packet and a **NEW unopened** lancet.



NOTE: Each test can be used **ONLY ONE TIME**. Do not try to use the test more than once.









Prepared on December 22, 2009. V1.0. Since manufacturers' instructions may have changed after this job aid was produced, all details should be cross-checked against manufacturer instructions in the product insert of the test in use.